

Monopole condensates in SU(N) Yang-Mills Theory

Vipul Periwal

Department of Physics, Princeton University, Princeton, New Jersey 08544

Faddeev and Niemi have proposed a reformulation of SU(2) Yang-Mills theory in terms of new variables, appropriate for describing the theory in its infrared limit based on the intuitive picture of colour confinement due to monopole condensation. I generalize their proposal (with some differences) to SU(N) Yang-Mills theory. The natural variables are $N - 1$ mutually commuting traceless $N \times N$ Hermitian matrices, an element of the maximal torus defined by these commuting matrices, $N - 1$ Abelian gauge fields for the maximal torus gauge group, and an invariant symmetric two-index tensor on the tangent space of the maximal torus, adding up to the requisite $2(N^2 - 1)$ physical degrees of freedom.

Twenty-five years after the discovery of asymptotic freedom [1], there is still no quantitative theoretical understanding of colour confinement in Yang-Mills theories. The qualitative picture of confinement [2] due to monopole condensation has yet to be given a quantitative justification starting from the fundamental Yang-Mills action.

Faddeev and Niemi [3] have recently proposed that even though the gauge field A_μ is the proper order parameter for describing the theory in its ultraviolet limit, in the infrared limit with monopole condensation some other order parameter may be more appropriate. They argue that in the infrared limit, for the SU(2) gauge theory, the appropriate order parameter is a three component vector $n^a(x)$ ($a = 1, 2, 3$) with unit length $\mathbf{n} \cdot \mathbf{n} = 1$ and classical action [4]

$$S = \int dx \left[m^2 (\partial_\mu \mathbf{n})^2 + \frac{1}{e^2} (\mathbf{n}, \mathbf{dn} \times \mathbf{dn})^2 \right] \quad (1)$$

with m a mass scale and e a dimensionless coupling constant. This is the unique local and Lorentz-invariant action for the unit vector $\mathbf{n}$ which is at most quadratic in time derivatives so that it admits a Hamiltonian interpretation, and involves all infra-red relevant (or marginal) terms with this property.

Following some ideas of Cho [5], they derived eq. 1 by parametrizing a generic four dimensional SU(2) connection

$$\mathbf{A}_\mu \equiv C_\mu \mathbf{n} + \mathbf{dn} \times \mathbf{n} + \rho \mathbf{dn} + \sigma \mathbf{dn} \times \mathbf{n}, \quad (2)$$

with an Abelian gauge field C_μ and scalar fields ρ, σ . They motivated this decomposition by considering the gauge transformation properties of A_μ under gauge transformations generated by $\mathbf{n}$, regarded as an element of the SU(2) Lie algebra. They noted that while A has $2 \cdot \dim(\text{SU}(2))$ physical degrees of freedom in four dimensions, C has two, $\mathbf{n}$ has two, and ρ, σ provide the requisite remaining two. In particular, an amusing suggestion [3] was that $\phi = \rho + i\sigma$ transforms as a complex scalar field under U(1) gauge transformations generated by $\alpha \cdot \mathbf{n}$. Note that this decomposition is specific to gauge theories in four dimensions. The effective description (eq. 1) is therefore dimension-specific—in other words, the action given in eq. 1 can be defined in any number of dimensions, but can describe low-energy Yang-Mills theory only in four dimensions.

Much intuition for the behaviour of gauge theories is based on the large N limit. It is therefore of interest to see how the Faddeev–Niemi decomposition generalizes for arbitrary gauge groups. For instance, what replaces the complex scalar field in the Faddeev–Niemi decomposition? Which degrees of freedom dominate in the large N limit? Cho already considered the SU(3) case [6] to some extent, but his arguments seem to involve an extension of Yang-Mills theory, and are therefore not directly related to the approach taken by Faddeev and Niemi [3]. I shall present below the general case of SU(N), which turns out to have considerable geometric intricacy compared to SU(2). It appears as somewhat of a miracle that the general parametrization given below (eq. 12) actually leads to a correct matching of degrees of freedom for any N (eq. 5). The final form will not exactly match the properties remarked on by Faddeev and Niemi [3] for SU(2), in that the variables for arbitrary N do not appear to admit a straightforward Higgs interpretation.

For SU(N), it is obvious that $\mathbf{n}(x)$ is replaced by $N - 1$ commuting traceless Hermitian $N \times N$ matrices, $H_I(x)$, orthonormal with respect to the trace, $\text{tr} H_I H_J = \delta_{IJ}$. How many parameters does it take to specify such a set of matrices? The simplest count is to consider that the dimension of the space of maximal tori of SU(N) is $N^2 - N$, and we further need to specify an orthonormal frame in the tangent space of the maximal torus, which requires an SO($N-1$) transformation, giving $N^2 - N + (N - 2)(N - 1)/2$ degrees of freedom. We thus have an $N - 1$ dimensional bundle of commuting, trace-orthonormal, traceless Hermitian matrices over spacetime. On this bundle, there is a natural spin connection, $\mathcal{D}_\mu H_I \equiv \partial_\mu H_I + \omega_{\mu I}{}^J H_J$, with ω specified completely by

$$\text{tr} H_I \mathcal{D}_\mu H_J = 0 \Rightarrow \omega_{\mu IJ} = -\text{tr} H_J \partial_\mu H_I. \quad (3)$$

The I, J indices on ω are raised and lowered with the identity matrix. $\omega_{\mu IJ}$ is antisymmetric in I, J since $\partial_\mu \text{tr} H_I H_J = 0$.

The point of introducing the spin connection (eq. 3) is that in terms of $\{H_I\}$, and $\mathcal{D}_\mu$, I can write

$$A_\mu \equiv C_\mu^I H_I + \phi^I \mathcal{D}_\mu H_I + i\beta^{IJ} [H_I, \mathcal{D}_\mu H_J] + \text{nonlinear terms}, \quad (4)$$

with $C_\mu^I \equiv \text{tr} A_\mu H_I$. Notice that $[H_I, \mathcal{D}_\mu H_J]$ is symmetric in I and J , since $\partial_\mu [H_I, H_J] = 0$. Thus $\mathcal{D}_\mu H_J$ and $[H_I, \mathcal{D}_\mu H_J]$ are linearly independent. For $\text{SU}(N)$, $N > 2$, there is no way to express further commutators of the form $[H_I, [H_K, \dots \mathcal{D}_\mu H_J] \dots]$ in terms of these two types of terms. As is obvious, such linearly independent higher commutators will automatically be generated by the action of gauge transformations. As I shall show, it is possible to include terms nonlinear in ϕ in such a way that gauge transformations corresponding to the maximal torus of $\text{SU}(N)$ defined by H_I preserve the complete form of A_μ .

Before considering the form of the requisite nonlinear terms, we should count degrees of freedom introduced in eq. 4. Besides the degrees of freedom in H_I , we get $2(N-1)$ from the $N-1$ Abelian gauge fields C_μ^I , $N-1$ from the scalar fields ϕ^I , and $N(N-1)/2$ from the scalar fields β^{IJ} . Therefore

$$2(N^2 - 1) = (N^2 - N + (N-2)(N-1)/2) + 2(N-1) + N-1 + N(N-1)/2, \quad (5)$$

which is exactly correct. In particular, if we added further degrees of freedom associated with multiple commutators, we would have too many degrees of freedom.

The appearance of the spin connection $\mathcal{D}_\mu$ leads to several significant differences from the $\text{SU}(2)$ case—obviously, the spin connection reduces to the ordinary derivative for the case of a real line bundle. Consider a gauge transformation generated by $\chi \equiv \alpha^I(x) H_I(x)$. Such gauge transformations are particularly significant in our rewriting of the gauge field, since they lie in the $\text{U}(1)^{N-1}$ maximal torus subgroup of $\text{SU}(N)$ gauge group defined by H_I . We have

$$\delta A_\mu = \partial_\mu \chi + i[A_\mu, \chi], \quad (6)$$

which we compare to

$$\delta A_\mu = \delta C_\mu^I H_I + \delta \phi^I \mathcal{D}_\mu H_I + i\delta \beta^{IJ} [H_I, \mathcal{D}_\mu H_J] + \delta(\text{nonlinear terms}). \quad (7)$$

Defining the action of the spin connection on α^I in the obvious manner, $(\mathcal{D}_\mu \alpha)^I \equiv \partial_\mu \alpha^I - \alpha^J \omega_{\mu J}^I$, we have

$$\partial_\mu \chi = (\mathcal{D}_\mu \alpha)^I H_I + \alpha^I (\mathcal{D}_\mu H)_I, \quad (8)$$

so that the Abelian gauge transformations of C_μ^I involve the spin connection:

$$\delta C_\mu^I = (\mathcal{D}_\mu \alpha)^I, \quad (9)$$

which is natural from the perspective of monopole moduli.

The behaviour of ϕ and β under such gauge transformations, eq. 6, is

$$\delta \phi^I = \alpha^I, \quad (10)$$

with

$$\delta \beta^{IJ} = -\frac{1}{2} (\phi^I \alpha^J + \phi^J \alpha^I). \quad (11)$$

However, we clearly need to find appropriate nonlinear terms to complete eq. 4, since not all terms in eq. 6 have been correlated with terms in eq. 7. The first step in finding these nonlinear terms is to note that eq. 11 shows that if we shift β by $\frac{1}{2} \phi^I \phi^J$, then the redefined β does not transform at all. Further, eq. 10 suggests that we should rewrite eq. 4 in a manner that makes this translation symmetry manifest.

Define $\text{Ad}_\phi(X) \equiv e^{-i\phi^I H_I} X e^{i\phi^I H_I}$. Consider then

$$A_\mu \equiv C_\mu^I H_I + \text{Ad}_\phi(E_\mu) - E_\mu + i\beta^{IJ} [H_I, \text{Ad}_\phi(\mathcal{D}_\mu H_J)], \quad (12)$$

where E_μ is implicitly defined by $-i[H_I, E_\mu] \equiv \mathcal{D}_\mu H_I$. (Notice that eq. 12 can be computed without ever computing E_μ explicitly, so the introduction of E_μ is solely to make eq. 12 transparent. However, the intuition for writing eq. 12 in terms of E_μ comes from observing that $[H_I, \mathcal{D}_\mu H_J]$ is symmetric in I and J , and that $\text{tr} H_I \mathcal{D}_\mu H_J = 0$.) An alternate way of writing eq. 12 is as

$$A_\mu + E_\mu = \text{Ad}_\phi(C_\mu^I H_I + i\beta^{IJ} [H_I, \mathcal{D}_\mu H_J] + E_\mu). \quad (13)$$

We deduce from eq. 12 that ϕ^I should be regarded as compact variables taking values on the maximal torus defined by H_I —this should be compared to the variables introduced in [3].

Now, under an infinitesimal gauge transformation with parameter $\alpha \cdot H \equiv \alpha^I H_I$, we have

$$A_\mu(C_\mu, \phi, \beta) \rightarrow A'_\mu = A_\mu(C_\mu + \mathcal{D}_\mu \alpha, \phi + \alpha, \beta). \quad (14)$$

This remarkably simple form, eq. 14, implies that under a gauge transformation with parameter $\alpha^I = -\phi^I$, the transformed gauge connection takes the form

$$A'_\mu = (C_\mu - \mathcal{D}_\mu \phi) \cdot H + i\beta^{IJ} [H_I, \mathcal{D}_\mu H_J] + \text{terms nonlinear in } \phi, \quad (15)$$

which makes the provenance of the physical degrees of freedom more transparent. The nonlinear terms are

$$E_\mu - \int_0^1 ds \text{Ad}_{-s\phi}(\phi \cdot \mathcal{D}_\mu H) - \text{Ad}_{-\phi}(E_\mu). \quad (16)$$

There is, of course, no gauge symmetry evident in eq. 15, since the field-dependent gauge transformation has fixed it completely, leading to a gauge in which A' contains no terms linear in ϕ , other than the derivative term appearing with C_μ .

What is the analogue of eq. 1 for $\text{SU}(N)$? Computing the curvature of the $\text{SU}(N)$ connection, as parametrized in eq. 12, and inserting into the Yang-Mills action, neglecting terms involving ϕ and C for the nonce, with $\beta^{IJ}(x) = \lambda \delta^{IJ} = \text{const.}$, we find

$$S = - \int dx \text{tr} (2\lambda [\partial_\mu H_I, \partial_\nu H^I] - \lambda^2 [[H_I, \partial_\mu H^I], [H_K, \partial_\nu H^K]])^2. \quad (17)$$

Notice that $\lambda = 0$ is a singular point, which is entirely appropriate when we recall the motivation of Cho [5,6], and Faddeev and Niemi [3]— $\beta = 0$ amounts essentially to enforcing that the monopole condensate does not contribute to the gauge potential in eq. 12. It should also be noted that local frame rotations $H_I(x) \rightarrow O_I^J(x) H_J(x)$ are *not* a symmetry of the action, but physical degrees of freedom (which have been included in eq. 5). While this action (eq. 17) admits a Hamiltonian interpretation, and also possesses infrared relevant deformations that would not appear in the Yang-Mills action, for example,

$$\Delta S = \int dx \{m_1^2 \text{tr} \partial_\mu H \cdot \partial^\mu H + m_2^2 \text{tr} [H_I, \partial_\mu H_K] [H^I, \partial^\mu H^K]\}, \quad (18)$$

it is absolutely crucial to keep in mind that the ϕ^I variables live on a maximal torus, related by a gauge invariance to the C_μ fields, so it is not correct to neglect these fields. A relevant deformation of the form

$$S' = \int dx M^2 (C_\mu - D_\mu \phi)^2 \quad (19)$$

preserves (Abelian) gauge invariance (eq. 14), and would serve to render the the gauge field massive (and hence eliminate it from the low energy dynamics of H if M were large enough relative to m_i). As required, the ϕ^I fields carry exactly enough degrees of freedom to provide all three polarizations of massive vector bosons, when combined with C_μ^I . In view of the fact that we are attempting a description of the confining behaviour of Yang-Mills theory, it is extremely important that *all* excitations have the *possibility* of acquiring a mass gap.

I do not go into more detail here regarding eq. 17 or its relevant perturbations because this is properly a matter of detailed calculation. The entirely reasonable claim [3] is that these relevant operators will appear when one integrates out higher momentum modes in a Wilsonian approach. In fact, one expects that all the relative coefficients of all the relevant perturbations should be calculable in such an approach. (There are other relevant operators besides those mentioned above.) I hope to return to this elsewhere.

Comparing eq. 12 to the form (eq. 2) suggested by Faddeev and Niemi [3], $\rho = (1 - \beta) \sin \phi$, and $\sigma = (1 - \beta) \cos \phi$. β does not transform under the Abelian gauge symmetry, hence an expectation value for β does not give rise in an obvious manner to a Higgs-like phase. However, since $(1 - \beta)^2 = \rho^2 + \sigma^2$, roughly speaking $1 - \beta$ is the magnitude of the Higgs field. So $\beta = 1$ is the phase with vanishing Higgs expectation value, which is also the value at which the Lagrangian for H simplifies. On the other hand, when $\beta = 0$, the Lagrangian for the H variables vanishes, which

one could call a Higgs phase, but is so singular that one has to be a trifle more careful in determining exactly what happens in the dynamics.

In conclusion, I have found a simple generalization of the decomposition suggested by Faddeev and Niemi [3] to arbitrary rank gauge groups. The natural variables in the $SU(N)$ case are H_I , a maximal Abelian subalgebra of the Lie algebra, ϕ^I , which are essentially coordinates on the maximal torus defined by H_I , β^{IJ} , a symmetric two-index tensor field on the tangent space of this maximal torus, and C_μ^I , $N-1$ Abelian gauge fields associated with H_I . Rather miraculously, the physical degrees of freedom in the $SU(N)$ gauge field match the physical degrees of freedom in these fields, eq. 5. The H_I and β^{IJ} fields have $O(N^2)$ degrees of freedom and may be expected to dominate the dynamics in the large N limit. From a canonical point of view, this change of variables is perfectly benign in the classical Hamiltonian. From the quantum point of view, the construction of a confinement vacuum may be simpler in the new variables. I have not addressed any issues of electric-magnetic duality for the simple reason that it is not at all clear that the change of variables (eq. 12) *should* make both the confinement and the Higgs phases of Yang-Mills theory transparent [7]. Indeed, one might expect that an electric version of eq. 12 would be a more appropriate starting point for describing a possible Higgs phase. A low-energy effective action, governing the dynamics of the H_I field is calculable starting from the Yang-Mills action and eq. 12, and should possess knot-like solutions analogous to those found by Faddeev and Niemi [8]. Further, it would be interesting to compare this action to [9].

Acknowledgements I am grateful to Wati Taylor and Peter Schupp for helpful comments. I am especially grateful to Y.J. Ng in correcting my interpretation of [3]. This work was supported in part by NSF grant PHY96-00258.

-
- [1] D.J. Gross and F. Wilczek, Phys. Rev. Lett. **30** 1343 (1973); H.D. Politzer, Phys. Rev. Lett. **30** 1346 (1973); G. 't Hooft, unpublished
 - [2] Y. Nambu, Phys. Rev. **D10** 4262 (1974); S. Mandelstam, Phys. Lett. **53B** 476 (1975); Phys. Rep. **23C** 245 (1976); Phys. Rev. **D19** 2391 (1978); G. 't Hooft, Nucl. Phys. **B138** 1 (1978); **B153** 141 (1979); in *High Energy Physics*, Proc. Euro. Phys. Soc. Int. Conf., Palermo, 1975, ed. A. Zichichi (Editrice Compositori, Bologna, 1976), p. 1225
 - [3] L.D. Faddeev and A.J. Niemi, hep-th/9807069; see also M. Mathur, hep-th/9807167
 - [4] L.D. Faddeev, IAS preprint IAS-75-QS70 (1970)
 - [5] Y.M. Cho, Phys. Rev. **D21** 1080 (1980); **D23** 2415 (1981)
 - [6] Y.M. Cho, Phys. Rev. Lett. **46** 302 (1981)
 - [7] S. Mandelstam, Phys. Rev. **D19** 2391 (1978)
 - [8] L.D. Faddeev and A.J. Niemi, Nature **387** 58 (1997), hep-th/9705176
 - [9] K. Bardakci and S. Samuel, Phys. Rev. **D18** 2849 (1978)